

CLEMSON SPINELESS

Okra

OUTSTANDING QUALITIES

- ◆ EASY HARVESTING
- ◆ HIGH YIELD POTENTIAL
- ◆ UNIFORM ATTRACTIVE FRUIT

Clemson Spineless is a main season standard Okra with very wide adaptability. Fruit are uniform, attractive green and have a good shelf life. Fruit have a good flavour and is almost spine free improving the ease of picking. Plants are strong and vigorous but easy to harvest. **Clemson Spineless** is widely adapted. Suited for both fresh and export market.



SPECIAL VARIETAL REQUIREMENTS

- None

CHARACTERISTIC*	CLEMSON SPINELESS
KIND	Okra (<i>Abelmoschus esculentus</i>)
PRODUCTION SYSTEM	Open field
MATURITY	Very early (approx. 55 - 65 days from sowing)
FRUIT DIMENSIONS	Usually harvested between 50 - 100 x 10 - 20 mm
FRUIT SHAPE	Pentagonal, tapered
LEAF SIZE	18 - 20 cm
SPINE	Almost spine free
EXTERIOR COLOUR	Bright, medium green
FLAVOUR	Good
VIGOUR	High
PLANT	Tall, half spreading, medium leafed and around 1.2 – 1.5 m plant height
BEARING HABIT	Upright
UNIFORMITY	Very good
OPTIMAL GROWTH TEMPERATURE	22 – 30 °C
AVERAGE SEED COUNT	160 - 185 seeds per g
POPULATION GUIDE	16 000 - 20 000 final stand per ha (60 cm in rows by 90 cm between rows)
USE	Fresh market, pre-pack, export
SPECIAL FEATURES	High yielding and widely adapted

* Characteristics given are affected by production methods such as soil type, nutrition, planting population, planting date and climatic conditions. Please read disclaimer.

Disclaimer: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Sakata Seed Southern Africa (Pty) Ltd Conditions of Sale before ordering seed.

Resistance: is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance, IR = Intermediate resistance).

Recent version: Kindly contact Sakata or Area Representative for the most recent version of this Technical Bulletin.